



Causes of variation in wild bee responses to anthropogenic drivers

Daniel P Cariveau¹ and Rachael Winfree

Anthropogenic change can have large impacts on wild bees and the pollination services they provide. However, the overall pattern of wild bee response to drivers such as land-use change, pesticides, pathogens, and climate change has been one of variability in both the magnitude and directionality of responses. We argue that two causes contribute to this variation. First, different species exhibit differential responses to the same anthropogenic drivers. Second, these anthropogenic drivers vary in type and magnitude that will drive variation in bee responses. For this second issue, we focus on land-use change, the most well-studied driver. We conclude by discussing how understanding species-level responses and the magnitude of land-use change can make bee conservation more effective.

Address

Department of Ecology, Evolution and Natural Resources, 14 College Farm Road, Rutgers University, New Brunswick, NJ 08901, USA

Corresponding author: Cariveau, Daniel P (dancariveau@gmail.com)

¹ Current address: Department of Entomology, University of Minnesota, 1980 Folwell Avenue, Falcon Heights, MN 55108, USA.

Current Opinion in Insect Science 2015, 10:104–109

This review comes from a themed issue on **Social insects**

Edited by **Christina Grozinger** and **Jay Evans**

<http://dx.doi.org/10.1016/j.cois.2015.05.004>

2214-5745/© 2015 Elsevier Inc. All rights reserved.

Introduction

Bees (Hymenoptera: Apoidea: Anthophila) are among the most ubiquitous and important pollinators of wild plants globally [1,2]. Furthermore, wild bee species, along with managed honey bees (*Apis mellifera*) and managed bumble bees (*Bombus* spp.), are important crop pollinators [3–5]. A few studies have shown regional [6] and local [7] declines in overall wild bee species richness. Declines in the best-studied genus, the bumble bees, is well-documented [8,9]. Given the ecological and agricultural importance of wild bees, these findings have raised concern among scientists, governments, and the general public. An array of anthropogenic drivers has been implicated in wild bee declines including pesticides, introduced pathogens,

climate change, and land-use change [10–12]. Furthermore, multiple interacting drivers may have even stronger negative effects [12,13]. However, bee responses to anthropogenic drivers are far from universal, showing a range of magnitudes and directionalities [14–16]. For example, a recent review of bee responses to the largest global driver of species loss, land-use change, found that while a 42% of effect sizes showed a negative response, 45% were neutral and 13% were positive [17].

The purpose of this essay is to explore the causes of variability in responses of bees to anthropogenic drivers. We have three main objectives. First, we highlight recent research that explores variability in species and species-group level responses of wild bees to anthropogenic drivers. Second, we examine how variation in an anthropogenic driver itself mediates wild bee responses. We focus on the most well-studied driver, human land-use change, which is also the leading cause of species loss globally [18], and describe how careful consideration of the magnitude and type of land use can reveal patterns of bee response that are otherwise obscured [17]. Third, we demonstrate how focusing on the responses of particular types of bee species can make conservation and management of wild bees more effective.


Variation in species-level responses

There are over 20 000 species of bees globally and they have a variety of life history traits and behaviors. It is therefore not surprising that different species show differential responses to the same anthropogenic drivers [19,20], and perhaps for this reason, also show different population trends over time [21**]. Recent species-specific studies move beyond simply assessing how aggregate wild bee abundance or species richness is affected by anthropogenic drivers (Figure 1), which had been the focus of the literature previously [17,22–24].

Pesticides

Pesticides are widely thought to be an important factor in wild bee declines [12], but bee species differ in their susceptibility to pesticides. A recent meta-analysis of lab-based, LD₅₀ studies examined 18 bee species other than honey bees and found that species exhibit differential susceptibilities to direct application of insecticides [20]. These differences in acute susceptibility may be due to species-level differences in body size, hemolymph chemistry, and

Figure 1



Schematic representing how focusing solely on total wild bee abundance obscures species-specific responses to an anthropogenic driver. Each solid, colored line represents the abundance of an individual bee species across a gradient of intensity for a hypothetical anthropogenic driver. The dashed line represents total bee abundance across that gradient.

immune system function [reviewed in 20]. Different classes of pesticides have different levels of toxicity across bee species [20]. For example, Biddinger *et al.* [25] examined mortality of honey bees and *Osmia cornifrons* (Megachilidae) to five different pesticides and found bee species by pesticide type interactions. For example, the LD₅₀ was met after applying 4 µg/bee of Acetamiprid to *O. cornifrons* compared to 65 µg/bee for honey bees but for Imidicloprid the LD₅₀ was 3.8 µg/bee for *O. cornifrons* compared to 0.2 µg/bee for honey bees. The recent literature is increasingly finding sublethal effects as well, particularly for the systemic neonicotinoids [26]. Given the ubiquity of these insecticides, any species-level variation in susceptibility to neonicotinoids could have substantial ecological effects.

Behavioral and natural history variation among bee species will likely make species-level variability in response to pesticides even greater in field settings than in the lab, although few field studies have explored this yet. Different behaviors and natural histories will result in different likelihoods of exposures among bee species. For example, bee species that have flight times that overlap with pesticide applications, and species whose host plants are concentrated in the area of application will be most susceptible [27]. For systemic pesticides such as neonicotinoids, pollen typically has higher concentrations than nectar [28,29] and thus may have differential effects on species that are foraging for either resource. Landscape-scale studies are crucial to predict which bee species will be most susceptible to pesticides. In one of few such studies, Rundlöf *et al.* [30••] compared bee responses in landscapes with and without neonicotinoid-treated crops

and found that bumble bee queen numbers and wild bee density was reduced with neonicotinoids while honey bee colony size showed no response.

Emerging pathogens from managed bees

The introduction of new diseases is a major concern for wild bees, with some species in decline likely due to pathogens, while others are unaffected. The best studied case concerns the bumble bee species in the subgenus *Bombus sensu stricto* which exhibited rapid declines in the midwestern United States [8]. This decline is associated with infection by the putatively introduced fungal pathogen *Nosema bombi*, which is commonly found in species in the subgenus *Bombus sensu stricto*. Spatially co-occurring species from other *Bombus* subgenera rarely host *N. bombi*, and these species are still relatively stable or increasing [8]. However, in the arctic and subarctic region of North America, some species in the subgenus *Bombus* show high *N. bombi* infection rates yet their populations are stable [31]. This example suggests that predicting which species will be most affected by emerging diseases will be challenging. Models with other insects suggest that diseases are most likely to affect species that are closely related to the hosts that harbor the new pathogens [32]. Thus given the ubiquity and global transport of honey bees, there is particular concern about their transmitting pathogens to others species of Apidae. Honey bees and bumble bees have been separated for over 70 million years [33], yet honey bee pathogens detrimentally infect bumble bees in laboratory settings [34–37] and vice versa [38•]. Honey bee pathogens have been found in multiple species of wild bees, but almost nothing is known about whether these pathogens have negative effects on wild species other than *Bombus* [39]. As domesticated bumble bees and honey bees are the most likely means by which these pathogens reach new locations, increased monitoring and control of these colonies is needed to protect wild bees [38•].

Climate change

Climate change will undoubtedly have differential effects on different bee species, as it does for other taxa [14]. A central focus of climate change research for bees has been potential asynchrony between bloom times and bee emergence [40]. Floral specialist bees could experience phenological mismatch differently from generalists. Two contrasting predictions have been made about this. First, some ecologists have predicted that specialist bees are at greater risk because if they emerge before or after their host plant blooms, they will be unable to forage [40]. However, some studies making this prediction are problematic as they confound rarity and sampling effects with true specialization [41]. Although there are well-known specialist (oligolectic) bee species [42], it is also true that species that appear to be specialists in a given study may not be specialists when more individuals are sampled, or when greater temporal and spatial scales are considered

[43]. Second, it has been proposed that specialist bees will be buffered against climate change because they have been selected to use the same cues for emergence that their host plants do; thus while these bees might change their phenology with climate warming, they would remain synchronized with their host plants [44,45^{*}]. Generalist species will likely be little effected by asynchrony as they can visit numerous plant species as has been demonstrated in some studies [46,47].

Physiological differences among bee species might also make them differentially affected by climate change. For example, as temperature increases, species that overwinter as adults lose more body mass during winter diapause which is likely due to greater metabolic rates and energy expenditure during winter [48]. This suggests that these species are more likely to negatively affected as the climate warms. In the same experiment, Fründ *et al.* [48] found that these effects were more pronounced in earlier emerging bees, suggesting that, as is the case for other taxa, early spring species will be the most affected by climate change [49].

Land-use change

Land-use change is one of the most important and best-studied anthropogenic drivers of bee declines, and response to land use varies strongly across bee species. One group of bee species, the floral specialist bees, consistently shows stronger declines with land-use change as compared to more generalist species. Often the decline of specialist species can be linked specifically to declines of their host plants [7,50,51,52^{**},53,54]. For example, in Europe, bee species that specialize on Fabaceae have experienced greater declines. This is due to changes in agricultural management, specifically a decline in planting Fabaceae as cover crops [52^{**}]. Specialist bees have shown to be disproportionately negatively affected in urban systems as well [53]. Conversely, populations of specialist bees can increase with land-use change if their host plants increase. In Europe, bee species that specialize on Rosaceae have increased along with increased plantings of Rosaceae crops [52^{**}]. Some species have likely expanded their range following plantings of their host plant [55]. Therefore, while specialist species as a group are more negatively affected by land-use change than generalists [51], these responses are species-specific and dependent on changes in host plant abundance.

By contrast, other traits associated with species or groups of species do not strongly predict response to land-use change. No consistent patterns have been found for body size [53,56], nesting guild [7,53], or sociality [50,57]. This may be due to the statistical difficulty of separating the causal role of traits that are correlated across taxa [51]. For example, the well-studied genus *Bombus* (the bumble bees) are social, large, and polylectic, making it more

difficult to determine which trait drives their responses to land-use change.

Variation in anthropogenic drivers: the case of land-use change

Another reason for the high variability in responses of bees to anthropogenic drivers is the variation in the anthropogenic drivers themselves. In this context, the relevant variation in responses includes not only that across bee species, but also in aggregate bee richness and abundance. We focus here on a single driver, land-use change, as this is the best studied of the anthropogenic drivers. Variation in land use has been found to influence the response of wild bee communities in two ways. First, the magnitude of the land-use change has strong effects on outcomes. For example, in systems where human land use is extreme (defined as some sites having $\leq 5\%$ of natural habitat remaining), bee abundance and richness decline steeply [17]. However in systems with less extreme land use (all sites have $> 5\%$ natural habitat), bee responses are mixed [17]. Importantly, in this review, 75% of the studies were done in systems with extreme human land use, which could bias the general impression that workers in this field have about how strongly bees are declining due to land-use change [17].

Second, bee responses vary with the type of land-use conversion studied. Bee abundance and species richness can increase with increasing urbanization and suburbanization [53,58]. Similarly, likely because many bee species are associated with open habitats, conversion of forest to more human-disturbed, open areas often increases bee abundance and richness [17]. Most studies of bee responses to land-use change have focused on conversion of natural habitat to agriculture, but within this broad category, different types of agriculture have differential effects [17]. For example in landscapes with red clover (*Trifolium pretense* L.), bumble bee queen densities were five time greater at sites with red clover than sites without [59]. By contrast to other studies that have demonstrated a negative effects of agriculture [e.g. 51].

Conserving wild bee communities

In this final section, we explore how the understanding of variability in species responses to anthropogenic drivers and variation in the drivers themselves can be applied to the conservation and management of wild bees. Understanding variation in species responses can provide valuable insight into an ongoing debate within conservation biology, which is: should conservation efforts focus on species that provide ecosystem services [61], or on the conservation of biodiversity per se, which generally emphasizes rare or threatened species [62]. A key question in this debate is whether the species that contribute ecosystem services such as pollination are also the important species driving patterns of biodiversity [63]. Recent studies suggest that only a small number of highly

abundant wild bee species do the great majority of crop pollination [64*,65]. By contrast, rare species make up most of the biodiversity [65]. Specifically, a large synthetic study across 90 different field studies found 785 bees species visit crops yet 80% of pollination was provided by only 2% of those bee species, which were the more common species [64*].

Given that the important crop pollinating species and rare bee species might often be two distinct sets of species, it is crucial to understand how each of these groups responds to anthropogenic drivers as well as conservation actions. Few studies have examined whether these different groups respond differently to a given anthropogenic driver. Important crop pollinating bees have been shown to persist in agricultural landscapes [64*]. However, we know of no studies that have directly measured how well rare bees, as a group, persist in agricultural landscapes. It may be that common, crop-pollinating species that persist in agricultural landscapes are more robust to pesticides as these chemicals are ubiquitous on many farms. In support of this hypothesis, Brittain *et al.* [66] found that species richness declined with pesticide application yet pollination was not affected, suggesting that biodiversity and ecosystem services are not responding similarly to this anthropogenic driver. It is also important to determine whether conservation actions benefit both important crop pollinators and rare bees. The common conservation action of planting of pollinator habitat may benefit both important crop pollinators and rare bees [67]. However, no studies have examined whether conservation efforts that optimize conservation of one group results in trade-offs in the conservation of the other.

The variation in anthropogenic drivers themselves can have important implications for designing conservation actions, in particular to determining the most effective locations for conservation action. For example, the effectiveness of pollinator habitat plantings varies with the magnitude of land-use change in the larger landscape surrounding the planted site. When pollinator habitat plantings are placed in landscapes with very intensive human land use (<1% surrounding landscape in natural habitat within a 1000 m radius) few bees use the habitat plantings, likely because the potential colonist species pool is highly reduced [68**]. By contrast, in landscapes with extensive natural habitat (>20% surrounding landscape in natural habitat within a 1000 m radius) pollinator habitat plantings have little positive effect on bee communities because these landscapes already support diverse and abundant bees [68**]. The most effective location for pollinator habitat plantings are landscapes with intermediate levels of human land use (1–20%). In these locations, pollinator habitat plantings significantly increase bee abundance and richness [68**].

Conclusions

As global change intensifies, there is an increasing potential for negative effects on wild bees. The effect of this change will depend on bee species, and on the magnitude and type of anthropogenic driver. We have made significant progress in understanding how wild bee abundance and richness, in aggregate, respond to land-use change. However, few studies have directly addressed the impact of pesticides, pathogens and climate change on multiple wild bee species in field settings [but see 30**]. In addition, no studies have quantified the relative effects of anthropogenic drivers across species. Doing so will enable for more effective prioritization of conservation and management efforts. Finally, wild bees must respond to multiple anthropogenic drivers simultaneously and this may increase negative effects [12,13]. These more nuanced approaches will lead to more accurate assessment of bee responses and allow for more effective conservation of these ecologically and economically important organisms.

Acknowledgements

We thank members of the Winfree lab for helpful comments. All photographs in Figure 1 are licensed under Creative Commons 3.0. Attributions from top to bottom are Alvesgaspar (*Bombus terrestris*), Alvesgaspar (*Anthidium florentinum*), Gideon Pisanty (*Megachile albisepta*), and Gideon Pisanty (*Megachile incerta*).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Neff JL, Simpson BB: **Bees, pollination systems and plant diversity.** In *Hymenoptera and Biodiversity*. Edited by LaSalle J, Gauld ID. CAB International; 1993:143-167.
 2. Ollerton J, Winfree R, Tarrant S: **How many flowering plants are pollinated by animals?** *Oikos* 2011, **120**:321-326.
 3. Garibaldi LA, Steffan-Dewenter I, Winfree R, Aizen MA, Bommarco R, Cunningham SA, Kremen C, Carvalheiro LG, Harder LD, Afik O *et al.*: **Wild pollinators enhance fruit set of crops regardless of honey bee abundance.** *Science* 2013, **339**:1608-1611.
 4. Klein AM, Vaissiere BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tscharntke T: **Importance of pollinators in changing landscapes for world crops.** *Proc R Soc B: Biol Sci* 2007, **274**:303-313.
 5. Velthuis HHW, van Doorn A: **A century of advances in bumblebee domestication and the economic and environmental aspects of its commercialization for pollination.** *Apidologie* 2006, **37**:421-451.
 6. Biesmeijer JC, Roberts SPM, Reemer M, Ohlemuller R, Edwards M, Peeters T, Schaffers AP, Potts SG, Kleukers R, Thomas CD *et al.*: **Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands.** *Science* 2006, **313**:351-354.
 7. Burkle LA, Marlin JC, Knight TM: **Plant-pollinator interactions over 120 years: loss of species, co-occurrence, and function.** *Science* 2013, **339**:1611-1615.
 8. Cameron SA, Lozier JD, Strange JP, Koch JB, Cordes N, Solter LF, Griswold TL: **Patterns of widespread decline in North American bumble bees.** *Proc Natl Acad Sci USA* 2011, **108**:662-667.

9. Schmid-Hempel R, Eckhardt M, Goulson D, Heinzmann D, Lange C, Plischuk S, Escudero LR, Salathe R, Scriven JJ, Schmid-Hempel P: **The invasion of southern South America by imported bumblebees and associated parasites.** *J Anim Ecol* 2014, **83**:823-837.
10. Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE: **Global pollinator declines: trends, impacts and drivers.** *Trends Ecol Evol* 2010, **25**:345-353.
11. Vanbergen AJ: **Initiative the insect pollinators: threats to an ecosystem service: pressures on pollinators.** *Front Ecol Environ* 2013, **11**:251-259.
12. Goulson D, Nicholls E, Botias C, Rotheray EL: **Bee declines driven by combined stress from parasites, pesticides, and lack of flowers.** *Science* 2015, **347**:1255-957.
13. Gonzalez-Varo JP, Biesmeijer JC, Bommarco R, Potts SG, Schweiger O, Smith HG, Steffan-Dewenter I, Szentgyorgyi H, Woyciechowski M, Vila M: **Combined effects of global change pressures on animal-mediated pollination.** *Trends Ecol Evol* 2013, **28**:524-530.
14. Bartomeus I, Park MG, Gibbs J, Danforth BN, Lakso AN, Winfree R: **Biodiversity ensures plant-pollinator phenological synchrony against climate change.** *Ecol Lett* 2013, **16**:1331-1338.
15. Rader R, Reilly J, Bartomeus I, Winfree R: **Native bees buffer the negative impact of climate warming on honey bee pollination of watermelon crops.** *Glob Change Biol* 2013, **19**:3103-3110.
16. Tuell JK, Isaacs R: **Community and species-specific responses of wild bees to insect pest control programs applied to a pollinator-dependent crop.** *J Econ Entomol* 2010, **103**:668-675.
17. Winfree R, Bartomeus I, Cariveau DP: **Native pollinators in anthropogenic habitats.** *Ann Rev Ecol Syst* 2011, **42**:1-22.
18. Pereira HM, Leadley PW, Proenca V, Alkemade R, Scharlemann JP, Fernandez-Manjarres JF, Araujo MB, Balvanera P, Biggs R, Cheung WW *et al.*: **Scenarios for global biodiversity in the 21st century.** *Science* 2010, **330**:1496-1501.
19. Cariveau DP, Williams NM, Benjamin FE, Winfree R: **Response diversity to land use occurs but does not consistently stabilise ecosystem services provided by native pollinators.** *Ecol Lett* 2013, **16**:903-911.
20. Arena M, Sgolastra F: **A meta-analysis comparing the sensitivity of bees to pesticides.** *Ecotoxicology* 2014, **23**:324-334.
21. Bartomeus I, Ascher JS, Gibbs J, Danforth BN, Wagner DL, Hedtke SM, Winfree R: **Historical changes in northeastern US bee pollinators related to shared ecological traits.** *Proc Natl Acad Sci USA* 2013, **110**:4656-4660.
- Using over 30 000 museum specimens collected from a 140-year period across the northeastern United States, the authors found that only 3 of 187 species experienced dramatic declines. However, 29% experienced modest declines in relative abundance and 27% of species experienced increases. Species-level traits were only weakly predictive of decline.
22. Wray JC, Neame LA, Elle E: **Floral resources, body size, and surrounding landscape influence bee community assemblages in oak-savannah fragments.** *Ecol Entomol* 2014, **39**:83-93.
23. Barlow J, Gardner TA, Araujo IS, Avila-Pires TC, Bonaldo AB, Costa JE, Esposito MC, Ferreira LV, Hawes J, Hernandez MI *et al.*: **Quantifying the biodiversity value of tropical primary, secondary, and plantation forests.** *Proc Natl Acad Sci USA* 2007, **104**:18555-18560.
24. Brosi BJ, Daily GC, Ehrlich PR: **Bee community shifts with landscape context in a tropical countryside.** *Ecol Appl* 2007, **17**:418-430.
25. Biddinger DJ, Robertson JL, Mullin C, Frazier J, Ashcraft SA, Rajotte EG, Joshi NK, Vaughn M: **Comparative toxicities and synergism of apple orchard pesticides to *Apis mellifera* (L.) and *Osmia cornifrons* (Radoszkowski).** *PLoS One* 2013, **8**:e72587.
26. Whitehorn PR, O'Connor S, Wackers FL, Goulson D: **Neonicotinoid pesticide reduces bumble bee colony growth and queen production.** *Science* 2012, **336**:351-352.
27. Brittain C, Potts SG: **The potential impacts of insecticides on the life-history traits of bees and the consequences for pollination.** *Basic Appl Ecol* 2011, **12**:321-331.
28. Dively GP, Kamel A: **Insecticide residues in pollen and nectar of a cucurbit crop and their potential exposure to pollinators.** *J Agric Food Chem* 2012, **60**:4449-4456.
29. Goulson D: **An overview of the environmental risks posed by neonicotinoid insecticides.** *J Appl Ecol* 2013, **50**:977-987.
30. Rundlöf M, Andersson GKS, Bommarco R, Fries I, Hederström V, Herbertsson L, Jonsson O, Klatt BK, Pedersen TR, Yourstone J *et al.*: **Seed coating with a neonicotinoid insecticide negatively affects wild bees.** *Nature* 2015, **521**:77-80.
- Wild bee density, nesting activity of a native bee (*Osmia bicornis* L.), and bumble bee colony development was reduced in landscapes containing oil seed rape treated with neonicotinoids. However, honey bee colony strength was not reduced suggesting variation in species responses to this driver. This study is a significant advance as the authors were able to compare bee responses among landscapes with and without neonicotinoid treatment.
31. Koch JB, Strange JP: **The status of *Bombus occidentalis* and *B. moderatus* in Alaska with special focus on *Nosema bombi* incidence.** *Northwest Sci* 2012, **86**:212-220.
32. Longdon B, Hadfield JD, Webster CL, Obbard DJ, Jiggins FM: **Host phylogeny determines viral persistence and replication in novel hosts.** *PLoS Pathog* 2011, **7**:e1002260.
33. Cardinal S, Danforth BN: **Bees diversified in the age of eudicots.** *Proc R Soc B: Biol Sci* 2013, **280**:20122686.
34. Graystock P, Yates K, Darvill B, Goulson D, Hughes WO: **Emerging dangers: deadly effects of an emergent parasite in a new pollinator host.** *J Invertebr Pathol* 2013, **114**:114-119.
35. Li J, Peng W, Wu J, Strange JP, Boncristiani H, Chen Y: **Cross-species infection of deformed wing virus poses a new threat to pollinator conservation.** *J Econ Entomol* 2011, **104**:732-739.
36. Meeus I, de Miranda JR, de Graaf DC, Wackers F, Smagge G: **Effect of oral infection with Kashmir bee virus and Israeli acute paralysis virus on bumblebee (*Bombus terrestris*) reproductive success.** *J Invertebr Pathol* 2014, **121**:64-69.
37. Fürst MA, McMahon DP, Osborne JL, Paxton RJ, Brown MJ: **Disease associations between honeybees and bumblebees as a threat to wild pollinators.** *Nature* 2014, **506**:364-366.
38. Graystock P, Yates K, Evison SEF, Darvill B, Goulson D, Hughes WOH, Osborne J: **The Trojan hives: pollinator pathogens, imported and distributed in bumblebee colonies.** *J Appl Ecol* 2013, **50**:1207-1215.
- Regulations in the UK are in place to limit the importation of bumble bee hives infected with parasites. Despite these regulations, 77% of hives were found to have microbial parasites with a number of these parasites demonstrating negative effects on bumble bee health.
39. Singh R, Levitt AL, Rajotte EG, Holmes EC, Ostiguy N, Vanengelsdorp D, Lipkin WI, Depamphilis CW, Toth AL, Cox-Foster DL: **RNA viruses in hymenopteran pollinators: evidence of inter-Taxa virus transmission via pollen and potential impact on non-*Apis* hymenopteran species.** *PLoS One* 2010, **5**:e14357.
40. Memmott J, Craze PG, Waser NM, Price MV: **Global warming and the disruption of plant-pollinator interactions.** *Ecol Lett* 2007, **10**:710-717.
41. Blüthgen N: **Why network analysis is often disconnected from community ecology: a critique and an ecologist's guide.** *Basic Appl Ecol* 2010, **11**:185-195.
42. Michener CD: *The Bees of the World*. 2nd edition. Johns Hopkins University Press; 2007.
43. Benadi G, Hovestadt T, Poethke HJ, Blüthgen N: **Specialization and phenological synchrony of plant-pollinator interactions along an altitudinal gradient.** *J Anim Ecol* 2014, **83**:639-650.

44. Forrest JRK, Thomson JD: **An examination of synchrony between insect emergence and flowering in Rocky Mountain meadows.** *Ecol Monogr* 2011, **81**:469-491.
45. Minckley RL, Roulston TH, Williams NM: **Resource assurance predicts specialist and generalist bee activity in drought.** *Proc R Soc B: Biol Sci* 2013, **280**:20122703.
- Bees and plants were sampled in a desert community were sampled before and after a drought. Bee species exhibited differential responses to the drought and these responses tracked those of their host plants.
46. Bartomeus I, Ascher JS, Wagner D, Danforth BN, Colla S, Kornbluth S, Winfree R: **Climate-associated phenological advances in bee pollinators and bee-pollinated plants.** *Proc Natl Acad Sci USA* 2011, **108**:20645-20649.
47. Forrest JRK: **Plant-pollinator interactions and phenological change: what can we learn about climate impacts from experiments and observations?** *Oikos* 2015, **124**:4-13.
48. Frund J, Zieger SL, Tschamtkke T: **Response diversity of wild bees to overwintering temperatures.** *Oecologia* 2013, **173**:1639-1648.
49. Parmesan C: **Ecological and evolutionary responses to recent climate change.** *Ann Rev Ecol Syst* 2006, **37**:637-669.
50. Rader R, Bartomeus I, Tylanakis JM, Laliberté E, van Kleunen M: **The winners and losers of land use intensification: pollinator community disassembly is non-random and alters functional diversity.** *Divers Distrib* 2014, **20**:908-917.
51. Williams NM, Crone EE, Roulston TH, Minckley RL, Packer L, Potts SG: **Ecological and life-history traits predict bee species responses to environmental disturbances.** *Biol Conserv* 2010, **143**:2280-2291.
52. Scheper J, Reemer M, van Kats R, Ozinga WA, van der Linden GT, Schaminee JH, Stepel H, Kleijn D: **Museum specimens reveal loss of pollen host plants as key factor driving wild bee decline in The Netherlands.** *Proc Natl Acad Sci USA* 2014, **111**:17552-17557.
- In this study, trends in species were determined using a national database with over 72 000 records across 98 years. Pollen loads were identified from museum species. The authors found that bee species that specialized on Fabaceae declined. This plant family has declined in the study region due to changes in farm management and loss of semi-natural habitat.
53. Cane JH, Minckley RL, Kervin LJ, Roulston TH, Williams NM: **Complex responses within a desert bee guild (Hymenoptera: Apiformes) to urban habitat fragmentation.** *Ecol Appl* 2006, **16**:632-644.
54. Bommarco R, Lundin O, Smith HG, Rundlöf M: **Drastic historic shifts in bumble-bee community composition in Sweden.** *Proc R Soc B-Biol Sci* 2012, **279**:309-315.
55. Bischoff I, Schröder S, Misof B: **Differentiation and range expansion of north american squash bee, *Peponapis pruinosa* (Apidae: Apiformes) populations assessed by geometric wing morphometry.** *Ann Entomol Soc Am* 2009, **102**:60-69.
56. Jauker B, Krauss J, Jauker F, Steffan-Dewenter I: **Linking life history traits to pollinator loss in fragmented calcareous grasslands.** *Landscape Ecol* 2013, **28**:107-120.
57. Winfree R, Aguilar R, Vazquez DP, LeBuhn G, Aizen MA: **A meta-analysis of bees' responses to anthropogenic disturbance.** *Ecology* 2009, **90**:2068-2076.
58. Matteson KC, Ascher JS, Langellotto GA: **Bee richness and abundance in New York city urban gardens.** *Ann Entomol Soc Am* 2008, **101**:140-150.
59. Rundlöf M, Persson AS, Smith HG, Bommarco R: **Late-season mass-flowering red clover increases bumble bee queen and male densities.** *Biol Conserv* 2014, **172**:138-145.
61. Kareiva P, Marvier M: **What is conservation science?** *BioScience* 2012, **62**:962-969.
62. Soule M: **The "new conservation".** *Conserv Biol* 2013, **27**:895-897.
63. Kremen C: **Managing ecosystem services: what do we need to know about their ecology?** *Ecol Lett* 2005, **8**:468-479.
64. Kleijn D, Winfree R, Bartomeus I, Carvalheiro LG, Henry M, Isaacs R, Klein A, Kremen C, M'Gonigle LK, Rader R, et al.: **Delivery of crop pollination services is an insufficient argument for wild pollinator conservation.** *Nat Commun* in press.
- Using a dataset of 90 different studies across the globe, the authors found that 2% of wild bee species found on crops contribute 80% of total pollination contributed to wild bees. In addition, these common, abundant species were relatively robust to land-use change.
65. Winfree R, Fox JW, Williams N, Reilly JR, Cariveau DP: **Abundance of common species, not species richness, drives delivery of a real-world ecosystem service.** *Ecol Lett* in press. <http://onlinelibrary.wiley.com/doi/10.1111/ele.12424/abstract>
66. Brittain C, Bommarco R, Vighi M, Barmaz S, Settele J, Potts SG: **The impact of an insecticide on insect flower visitation and pollination in an agricultural landscape.** *Agric Forest Entomol* 2010, **12**:259-266.
67. Pywell RF, Meek WR, Hulmes L, Hulmes S, James KL, Nowakowski M, Carvell C: **Management to enhance pollen and nectar resources for bumblebees and butterflies within intensively farmed landscapes.** *J Insect Conserv* 2011, **15**:853-864.
68. Scheper J, Holzschuh A, Kuussaari M, Potts SG, Rundlöf M, Smith HG, Kleijn D: **Environmental factors driving the effectiveness of European agri-environmental measures in mitigating pollinator loss — a meta-analysis.** *Ecol Lett* 2013, **16**:912-920.
- This meta-analysis tested whether landscape context influenced wild bee responses to agri-environment schemes across Europe. Wild bee abundance and richness exhibited the greatest increase when agri-environment schemes took place in simple as compared to highly cleared or complex landscapes. In addition, the benefit of the agri-environment schemes increased as the contrast between floral resources in agri-environment schemes the surrounding landscape increased.